

Contribution to the Proceedings of *Texas in Tuscany: XXI Symposium on Relativistic Astrophysics*, 9–13 December 2002, Florence, Italy.

NEUTRINOS IN PHYSICS AND ASTROPHYSICS

GEORG G. RAFFELT

*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut)
Föhringer Ring 6, 80805 München, Germany
E-mail: raffelt@mppmu.mpg.de*

The observed flavor oscillations of solar and atmospheric neutrinos determine several elements of the leptonic mixing matrix, but leave open the small mixing angle Θ_{13} , a possible CP-violating phase, the mass ordering, the absolute mass scale m_ν , and the Dirac vs. Majorana property. Progress will be made by long-baseline, tritium endpoint, and 2β decay experiments. The best constraint on m_ν obtains from cosmological precision observables, implying that neutrinos contribute very little to the dark matter. However, massive Majorana neutrinos may well be responsible for ordinary matter by virtue of the leptogenesis mechanism for creating the baryon asymmetry of the universe. In future, neutrinos could play an important role as astrophysical messengers if point sources are discovered in high-energy neutrino telescopes. In the low-energy range, a high-statistics observation of a galactic supernova would allow one to observe directly the dynamics of stellar collapse and perhaps to discriminate between certain mixing scenarios. An observation of the relic neutrinos from all past supernovae has come within reach.

1. Flavor Oscillations

Neutrino oscillations are now firmly established by measurements of solar and atmospheric neutrinos and the KamLAND and K2K long-baseline experiments^{1,2,3,4}. Evidently the weak interaction eigenstates ν_e , ν_μ and ν_τ are non-trivial superpositions of three mass eigenstates ν_1 , ν_2 and ν_3 ,

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}. \quad (1)$$

The leptonic mixing matrix can be written in the canonical form

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & e^{i\delta}s_{13} \\ 0 & 1 & 0 \\ -e^{-i\delta}s_{13} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (2)$$

where $c_{12} = \cos \Theta_{12}$ and $s_{12} = \sin \Theta_{12}$ with Θ_{12} the 12-mixing angle, and so forth. One peculiarity of 3-flavor mixing beyond the 2-flavor case is a

non-trivial phase δ that can lead to CP-violating effects, i.e. the 3-flavor oscillation pattern of neutrinos can differ from that of anti-neutrinos.

The atmospheric neutrino oscillations essentially decouple from the solar ones and are controlled by the 23-mixing that may well be maximal (45°). The solar case is dominated by 12-mixing that is large but not maximal. The Chooz reactor experiment provides an upper limit on the small 13-mixing. From a global 3-flavor analysis of all data one finds the 3σ ranges for the mass differences and mixing angles summarized in Table 1.

Table 1. Neutrino mixing parameters from a global analysis of all experiments¹ (3σ ranges).

Combination	Mixing angle Θ	Δm^2 [meV ²]
12	27° – 42°	55–190
23	32° – 60°	1400–6000
13	$< 14^\circ$	$\approx \Delta m_{23}^2$

The only evidence for flavor conversions that is inconsistent with this picture comes from LSND, a short-baseline accelerator experiment. If the excess $\bar{\nu}_e$ counts are interpreted in terms of $\bar{\nu}_\mu$ - $\bar{\nu}_e$ -oscillations, the allowed mixing parameters populate two islands within⁵ $\Delta m^2 = 0.2$ – 7 eV² and $\sin^2 2\Theta = (0.3$ – $5) \times 10^{-2}$. One possibility to accommodate this Δm^2 with the atmospheric and solar values is a fourth sterile neutrino appearing as an intermediate state to account for the LSND measurements, although this scheme is now almost certainly ruled out⁶. Another solution is the radical conjecture that the masses of neutrinos differ from those of anti-neutrinos, implying a violation of the CPT symmetry⁷.

Therefore, if LSND is confirmed by the ongoing MiniBooNE project⁸ the observed flavor conversions imply something far more fundamental than neutrino mixing. Many workers in neutrino physics take the attitude that this would be too good to be true and thus are skeptical about LSND. Either way, MiniBooNE is crucial to clarify this fundamental point.

Assuming MiniBooNE will refute LSND so that there is no new revolution, the mass and mixing parameters given in Table 1 still leave many questions open. Is the 23-mixing truly maximal while the 13-mixing is not? How large is the small 13-mixing angle? Is there a CP-violating phase? Moreover, it is possible that two mass eigenstates separated by the small “solar” mass difference could form a doublet separated by the large “atmospheric” difference from a lower-lying single state (“inverted hierarchy”).

These issues will be addressed by long-baseline experiments involving reactor and accelerator neutrinos. KamLAND and K2K in Japan are already taking data, while the Fermilab to Soudan and CERN to Gran Sasso projects, each with a baseline of 730 km, are under construction. Future projects involving novel technologies (superbeams, neutrino factories, beta-beams)^{9,10} and their physics potential^{11,12,13} are being discussed. The “holy grail” of these efforts is finding leptonic CP violation.

2. Neutrino Dark Matter and Cosmic Structure Formation

The most direct limit on the overall mass scale m_ν derives from tritium experiments searching for a deformation of the β end-point spectrum. The final limit from Mainz and Troitsk is¹⁴

$$m_\nu < 2.2 \text{ eV} \quad (95\% \text{ CL}). \quad (3)$$

This number is much larger than the mass splittings, obviating the need for a detailed interpretation in terms of mixed neutrinos. In future, the KATRIN experiment¹⁴ is expected to reach a sensitivity of 0.35 eV.

Traditionally it is cosmology that provides the most restrictive m_ν limits. Standard big bang cosmology predicts a present-day density of

$$n_{\nu\bar{\nu}} = \frac{3}{11} n_\gamma \approx 112 \text{ cm}^{-3} \quad \text{per flavor}. \quad (4)$$

This translates into a cosmic neutrino mass fraction of

$$\Omega_\nu h^2 = \sum_{i=1}^3 \frac{m_i}{92.5 \text{ eV}}, \quad (5)$$

where h is the Hubble parameter in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The oscillation experiments imply $m_\nu > 40 \text{ meV}$ for the largest neutrino mass eigenstate so that $\Omega_\nu > 0.8 \times 10^{-3}$ if $h = 0.72$. On the other hand, the tritium limit Eq. (3) implies $\Omega_\nu < 0.14$ so that neutrinos could still contribute significantly to the dark matter.

This possibility is severely constrained by large-scale structure observations. Neutrino free streaming in the early universe erases small scale density fluctuations so that the hot dark matter fraction is most effectively constrained by the small-scale power of the cosmic matter density fluctuations. The recent 2dF Galaxy Redshift Survey data imply^{15,16}

$$\sum m_\nu < 1.0 \text{ eV} \quad (95\% \text{ CL}). \quad (6)$$

To arrive at this limit other cosmological data were used, notably the angular power spectrum of cosmic microwave background radiation as measured

by WMAP as well as reasonable priors on other parameters such as the Hubble constant. If one includes the more problematic Lyman- α forest data to constrain the small-scale power of the matter density the limit improves to¹⁷ $\sum m_\nu < 0.69$ eV. The dependence of such limits on priors and other assumptions is discussed in the cited papers^{15,16}.

In future the Sloan Digital Sky Survey will improve the galaxy correlation function while additional CMBR data from WMAP and later from Planck will improve the matter power spectrum, enhancing the cosmological m_ν sensitivity^{18,19}. Especially promising are future weak lensing data^{20,21} that may come surprisingly close to the lower limit $\sum m_\nu > 40$ meV implied by the atmospheric neutrino oscillations.

While the progress in precision cosmology has been impressive one should keep worrying about systematic effects that do not show up in statistical confidence levels. Even when the cosmological limits are nominally superior to near-future experimental sensitivities, there remains a paramount need for independent laboratory experiments.

3. How Many Neutrinos in the Universe?

To translate a laboratory m_ν measurement or limit into a hot dark matter fraction Ω_ν and the reverse one usually assumes the standard cosmic neutrino density Eq. (4). However, thermal neutrinos in the early universe are characterized by unknown chemical potentials μ_ν or degeneracy parameters $\xi_\nu = \mu_\nu/T$ for each flavor. While the small baryon-to-photon ratio $\sim 10^{-9}$ suggests that all degeneracy parameters are small, large asymmetries between neutrinos and anti-neutrinos could exist and vastly enhance the overall density.

The recent WMAP measurement of the CMBR angular power spectrum provides new limits on the cosmic radiation density^{15,22,23}. However, the most restrictive limits on neutrino degeneracy parameters still obtain from big-bang nucleosynthesis (BBN) that is affected in two ways. First, a larger neutrino density increases the primordial expansion rate, thereby increasing the neutron-to-proton freeze-out ratio n/p and thus the cosmic helium abundance. Second, electron neutrinos modify $n/p \propto \exp(-\xi_{\nu_e})$. Depending on the sign of ξ_{ν_e} this effect can compensate for the expansion-rate effect of ν_μ or ν_τ so that no significant BBN limit on the overall neutrino density obtains²⁴. If ξ_{ν_e} is the only chemical potential, the observed helium abundance yields $-0.01 < \xi_{\nu_e} < 0.07$.

However, neutrino oscillations imply that the individual flavor lepton

numbers are not conserved so that in thermal equilibrium there is only one chemical potential for all flavors. If equilibrium is achieved before n/p freeze-out, the restrictive BBN limit on ξ_{ν_e} applies to all flavors, $|\xi_\nu| < 0.07$, fixing the cosmic neutrino density to within about 1%. The approach to flavor equilibrium by neutrino oscillations and collisions was recently studied^{25,26,27,28}. The details are subtle due to the large weak potential caused by the neutrinos themselves, causing the intriguing phenomenon of synchronized flavor oscillations^{29,30,31}.

The bottom line is that effective flavor equilibrium before n/p freeze-out is reliably achieved only if the solar oscillation parameters are in the favored LMA region. Now that KamLAND has confirmed LMA, for the first time BBN provides a reliable handle on the cosmic neutrino density. As a consequence, for the first time the relation between Ω_ν and m_ν is uniquely given by the standard expression Eq. (5).

4. Neutrino Majorana Masses and Leptogenesis

The neutrino contribution to the dark matter density is negligible. Intriguingly, however, they may play a crucial role for the baryon asymmetry of the universe (BAU) and thus the presence of ordinary matter³². The main ingredients of this leptogenesis scenario are those of the usual see-saw mechanism for small neutrino masses. The ordinary light neutrinos have right-handed partners with large Majorana masses. The left- and right-handed states are coupled by Dirac mass terms that obtain from Yukawa interactions with the Higgs field. The heavy Majorana neutrinos will be in thermal equilibrium in the early universe. When the temperature falls below their mass, their equilibrium density becomes exponentially suppressed. However, if at that time they are no longer in thermal equilibrium, their abundance will exceed the equilibrium distribution. The subsequent out-of-equilibrium decays can lead to the net generation of lepton number. CP-violation is possible by the usual interference of tree-level with one-loop diagrams. The generated lepton number excess will be re-processed by standard-model sphaleron effects which respect $B - L$ but violate $B + L$. It is straightforward to generate the observed BAU by this mechanism.

The requirement that the heavy Majorana neutrinos freeze out before they get Boltzmann suppressed implies an upper limit on the same parameter combination of Yukawa couplings and heavy Majorana masses that determines the observed small neutrino masses³³. Most recently, a robust

upper limit on all neutrino masses of

$$m_\nu < 120 \text{ meV} \quad (7)$$

was claimed³⁴. Degenerate neutrinos with a “large” common mass scale of, e.g., 400 meV require a very precise degeneracy of the heavy Majorana masses to better than 10^{-3} .

A necessary ingredient for this mechanism is the Majorana nature of neutrino masses that can be tested in the laboratory by searching for neutrinoless 2β decay. This process is sensitive to

$$\langle m_{ee} \rangle = \left| \sum_{i=1}^3 \lambda_i |U_{ei}|^2 m_i \right| \quad (8)$$

with λ_i a Majorana CP phase. Therefore, we have two additional physically relevant phases beyond the Dirac phase δ of the previously discussed mixing matrix. If neutrinos have Majorana masses their mixing involves three mass eigenstates, three mixing angles, and three physical phases.

Actually, several members of the Heidelberg-Moscow collaboration have claimed first evidence for this process^{35,36}, implying a 95% CL range of $\langle m_{ee} \rangle = 110\text{--}560$ meV. Uncertainties of the nuclear matrix element can widen this range by up to a factor of 2 in either direction. The significance of this discovery has been fiercely critiqued by many experimentalists working on other 2β projects³⁷. Even when taking the claimed evidence at face value the statistical significance is only about 97%, too weak for definitive conclusions. More sensitive experiments are needed and developed to explore this range of Majorana masses³⁸.

5. High-Energy Neutrinos From Astrophysical Sources

The 2002 Physics Nobel Prize was awarded, in part, to Raymond Davis and Masatoshi Koshiba “for pioneering contributions to astrophysics, in particular for the detection of cosmic neutrinos.” Unfortunately, the observed sources remain limited to the Sun and Supernova 1987A, apart from the diffuse flux of atmospheric neutrinos. This situation could radically change in the near future if the high-energy neutrino telescopes that are currently being developed begin to discover astrophysical point sources.

The spectrum of cosmic rays reaches to energies of at least 3×10^{20} eV, proving the existence of cosmic sources for particles with enormous energies^{39,40}. Most of the cosmic rays appear to be protons or nuclei so that there must be hadronic accelerators both within our galaxy and beyond. Wherever high-energy hadrons interact with matter or radiation, the

decay of secondary pions produces a large flux of neutrinos. At the source one expects a flavor composition of $\nu_e : \nu_\mu : \nu_\tau = 1 : 2 : 0$, but the observed oscillations imply equal fluxes of all flavors at Earth. High-energy neutrino astronomy offers a unique opportunity to detect the enigmatic sources of high-energy cosmic rays because neutrinos are neither absorbed by the cosmic photon backgrounds nor deflected by magnetic fields.

While there are many different models for possible neutrino sources^{40,41}, the required size for a detector is generically 1 km^3 . The largest existing neutrino telescope, the AMANDA ice Cherenkov detector at the South Pole, is about 1/10 this size. It has not yet observed a point source, but the detection of atmospheric neutrinos shows that this approach to measuring high-energy neutrinos works well⁴². It is expected that this instrument is upgraded to the full 1 km^3 size within the next few years under the name of IceCube⁴³. Similar instruments are being developed in the Mediterranean⁴⁴. Moreover, air-shower arrays for ordinary cosmic rays may detect very high-energy neutrinos by virtue of horizontal air showers⁴⁵. Although this field is in its infancy, it holds the promise of exciting astrophysical discoveries in the foreseeable future.

6. Supernova Neutrinos

The observation of neutrinos from the supernova (SN) 1987A in the Large Magellanic Cloud was a milestone for neutrino astronomy, but the total of about 20 events in the Kamiokande and IMB detectors was frustratingly sparse. The chances of observing a galactic SN are small because SNe are thought to occur with a rate of at most a few per century. On the other hand, many neutrino detectors and especially Super-Kamiokande have a rich physics program for perhaps decades to come, notably in the area of long-baseline oscillation experiments and proton decay searches. Likewise, the south pole detectors may be active for many decades and would be powerful SN observatories⁴⁶. Therefore, it remains worthwhile to study what can be learned from a high-statistics SN observation.

The explosion mechanism for core-collapse SNe remains unsettled as long as numerical simulations fail to reproduce robust explosions. A high-statistics neutrino observation is probably the only chance to watch the collapse and explosion dynamics directly and would allow one to verify the standard delayed explosion scenario⁴⁷. The neutrinos arrive a few hours before the optical explosion so that a neutrino observation can serve to alert the astronomical community, a task pursued by the Supernova Early Warn-

ing System (SNEWS)⁴⁸. For particle physics, many of the limits based on the SN 1987A neutrino signal⁴⁹ would improve and achieve a firm experimental and statistical basis. On the other hand, the time-of-flight sensitivity to the neutrino mass is in the range of a few eV,⁵⁰ not good enough as the “ m_ν frontier” has moved to the sub-eV range.

Can we learn something useful about neutrino mixing from a galactic SN observation? This issue has been addressed in many recent studies^{51,52,53,54}, and the answer is probably yes, depending on the detectors operating at the time of the SN, their geographic location, and the neutrino mixing scenario, i.e. the magnitude of the small mixing angle Θ_{13} and the ordering of the masses. Any observable oscillation effects depend on the spectral and flux differences between the different flavors. We have recently shown that previous studies overestimated these differences^{55,56,57} because traditional numerical simulations used a schematic treatment of ν_μ and ν_τ transport. Distinguishing, say, between the normal and inverted mass ordering remains a daunting task at long-baseline experiments. Therefore, a future galactic SN observation may still offer a unique opportunity to settling this question.

The relic flux from all past SNe in the universe is observable because it exceeds the atmospheric neutrino flux for energies below 30–40 MeV. Recently Super-Kamiokande has reported a limit that already caps the more optimistic predictions⁵⁸. Significant progress depends on suppressing the background caused by the decay of sub-Cherenkov muons from low-energy atmospheric neutrinos. One possibility is to include an efficient neutron absorber such as gadolinium in the detector that would tag the reactions $\bar{\nu}_e + p \rightarrow n + e^+$.⁵⁹ If this approach works in practice the detection of relic SN neutrinos has come within experimental reach.

7. Conclusions

After neutrino oscillations have been established, the next challenge is to pin down the as yet undetermined elements of the mixing matrix and the absolute masses and mass ordering. Long-baseline experiments can address many of these questions and may even discover leptonic CP-violation. The Majorana nature of neutrinos can be established in 2β experiments if the 0ν decay mode can be convincingly observed. Majorana neutrinos with masses < 120 meV fit nicely into the leptogenesis scenario for creating the baryon asymmetry of the universe so that neutrinos may well be responsible for the ordinary matter in the universe. Their contribution to the dark matter is non-zero but negligible. Still, precision observations of cos-

mological large-scale structure remain the most powerful tool to constrain the absolute mass scale, although independent laboratory confirmation remains crucial. In the past, neutrino oscillation physics was dominated by solar and atmospheric neutrinos, but long-baseline experiments are about to “take over.” In future, neutrinos from natural sources are likely more important as astrophysical messengers while oscillation physics will mostly be done in the laboratory. The search for astrophysical point sources with high-energy neutrino telescopes may soon open a new window to the universe. Observing a high-statistics neutrino signal from a future galactic supernova remains perhaps the most cherished prize for low-energy neutrino observatories. Meanwhile the search for the cosmic relic neutrinos from all past supernovae has become a realistic possibility. Neutrino physics and astrophysics will remain fascinating for a long time to come!

Acknowledgments

This work was supported, in part, by the Deutsche Forschungsgemeinschaft under grant No. SFB-375 and by the European Science Foundation (ESF) under the Network Grant No. 86 Neutrino Astrophysics.

References

1. M. C. Gonzalez-Garcia and Y. Nir, “Developments in neutrino physics,” hep-ph/0202058.
2. G. L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo and A. M. Rotunno, “Solar neutrino oscillation parameters after first KamLAND results,” hep-ph/0212127.
3. J. N. Bahcall, M. C. Gonzalez-Garcia and C. Pena-Garay, “Solar neutrinos before and after KamLAND,” *JHEP* **02**, 009 (2003) [hep-ph/0212147].
4. P. C. de Holanda and A. Y. Smirnov, “LMA MSW solution of the solar neutrino problem and first KamLAND results,” *JCAP* **02**, 001 (2003) [hep-ph/0212270].
5. E. D. Church, K. Eitel, G. B. Mills and M. Steidl, “Statistical analysis of different $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ searches,” *Phys. Rev. D* **66**, 013001 (2002) [hep-ex/0203023].
6. M. Maltoni, T. Schwetz, M. A. Tortola and J. W. Valle, “Global analysis of neutrino oscillation data in four-neutrino schemes,” *Nucl. Phys. Proc. Suppl.* **114**, 203 (2003) [hep-ph/0209368].
7. G. Barenboim, L. Borissov and J. Lykken, “CPT violating neutrinos in the light of KamLAND,” hep-ph/0212116.
8. A. O. Bazarko (BooNE Collaboration), “Status of MiniBooNE,” hep-ex/0210020.
9. Neutrino Factory & Muon Storage Rings at CERN <http://muonstoragerings.web.cern.ch/>

10. Neutrino Factory Working Group <http://nfwg.home.cern.ch/nfwg/>
11. V. D. Barger, S. Geer, R. Raja and K. Whisnant, "Exploring neutrino oscillations with superbeams," *Phys. Rev. D* **63**, 113011 (2001) [hep-ph/0012017].
12. A. Cervera, A. Donini, M. B. Gavela, J. J. Gomez Cadenas, P. Hernandez, O. Mena and S. Rigolin, "Golden measurements at a neutrino factory," *Nucl. Phys. B* **579**, 17 (2000); Erratum *ibid.* **593**, 731 (2001) [hep-ph/0002108].
13. M. Freund, P. Huber and M. Lindner, "Systematic exploration of the neutrino factory parameter space including errors and correlations," *Nucl. Phys. B*, **615**, 331 (2001) [hep-ph/0105071].
14. C. Weinheimer, "Direct neutrino mass search," hep-ex/0210050.
15. S. Hannestad, "Neutrino masses and the number of neutrino species from WMAP and 2dFGRS," astro-ph/0303076.
16. O. Elgaroy and O. Lahav, "The role of priors in deriving upper limits on neutrino masses from the 2dFGRS and WMAP," astro-ph/0303089.
17. D. N. Spergel *et al.*, "First year WMAP observations: Determination of cosmological parameters," astro-ph/0302209.
18. W. Hu, D. J. Eisenstein and M. Tegmark, "Weighing neutrinos with galaxy surveys," *Phys. Rev. Lett.* **80**, 5255 (1998) [astro-ph/9712057].
19. S. Hannestad, "Can cosmology detect hierarchical neutrino masses?," astro-ph/0211106.
20. W. Hu, "Dark energy and matter evolution from lensing tomography," *Phys. Rev. D* **66**, 083515 (2002) [astro-ph/0208093].
21. K. N. Abazajian and S. Dodelson, "Neutrino mass and dark energy from weak lensing," astro-ph/0212216.
22. P. Crotty, J. Lesgourgues and S. Pastor, "Measuring the cosmological background of relativistic particles with WMAP," astro-ph/0302337.
23. E. Pierpaoli, "Constraints on the cosmic neutrino background," astro-ph/0302465.
24. H. S. Kang and G. Steigman, "Cosmological constraints on neutrino degeneracy," *Nucl. Phys. B* **372**, 494 (1992).
25. C. Lunardini and A. Y. Smirnov, "High-energy neutrino conversion and the lepton asymmetry in the universe," *Phys. Rev. D* **64**, 073006 (2001) [hep-ph/0012056].
26. A. D. Dolgov, S. H. Hansen, S. Pastor, S. T. Petcov, G. G. Raffelt and D. V. Semikoz, "Cosmological bounds on neutrino degeneracy improved by flavor oscillations," *Nucl. Phys. B* **632**, 363 (2002) [hep-ph/0201287].
27. Y. Y. Wong, "Analytical treatment of neutrino asymmetry equilibration from flavour oscillations in the early universe," *Phys. Rev. D* **66**, 025015 (2002) [hep-ph/0203180].
28. K. N. Abazajian, J. F. Beacom and N. F. Bell, "Stringent constraints on cosmological neutrino antineutrino asymmetries from synchronized flavor transformation," *Phys. Rev. D* **66**, 013008 (2002) [astro-ph/0203442].
29. S. Samuel, "Neutrino oscillations in dense neutrino gases," *Phys. Rev. D* **48**, 1462 (1993).
30. S. Pastor, G. G. Raffelt and D. V. Semikoz, "Physics of synchronized neutrino oscillations caused by self-interactions," *Phys. Rev. D* **65**, 053011 (2002) [hep-

- ph/0109035].
31. S. Pastor and G. Raffelt, “Flavor oscillations in the supernova hot bubble region: Nonlinear effects of neutrino background,” *Phys. Rev. Lett.* **89**, 191101 (2002) [astro-ph/0207281].
 32. M. Fukugita and T. Yanagida, “Baryogenesis without grand unification,” *Phys. Lett. B* **174**, 45 (1986).
 33. W. Buchmüller and M. Plümacher, “Neutrino masses and the baryon asymmetry,” *Int. J. Mod. Phys. A* **15**, 5047 (2000) [hep-ph/0007176].
 34. W. Buchmüller, P. Di Bari and M. Plümacher, “The neutrino mass window for baryogenesis,” hep-ph/0302092.
 35. H. V. Klapdor-Kleingrothaus, A. Dietz, H. L. Harney and I. V. Krivosheina, “Evidence for neutrinoless double beta decay,” *Mod. Phys. Lett. A* **16**, 2409 (2001) [hep-ph/0201231].
 36. H. V. Klapdor-Kleingrothaus, “Neutrinoless double-beta decay: Status of evidence and future,” *Phys. Atom. Nucl.* **65**, 2135 (2002) [*Yad. Fiz.* **65**, 2198 (2002)].
 37. C. E. Aalseth *et al.*, “Comment on ‘Evidence for neutrinoless double beta decay’,” *Mod. Phys. Lett. A* **17**, 1475 (2002) [hep-ex/0202018].
 38. O. Cremonesi, “Neutrinoless double beta decay: Present and future,” hep-ex/0210007.
 39. G. Sigl, “Ultrahigh-energy cosmic rays: A probe of physics and astrophysics at extreme energies,” *Science* **291**, 73 (2001) [astro-ph/0104291].
 40. F. Halzen and D. Hooper, “High-energy neutrino astronomy: The cosmic ray connection,” *Rept. Prog. Phys.* **65**, 1025 (2002) [astro-ph/0204527].
 41. E. Waxman, “Astrophysical sources of high energy neutrinos,” astro-ph/0211358.
 42. J. Ahrens *et al.* (AMANDA Collaboration), “Search for point sources of high energy neutrinos with AMANDA,” *Astrophys. J.* **583**, 1040 (2003) [astro-ph/0208006].
 43. F. Halzen, “High-energy neutrino astronomy: Science and first results,” astro-ph/0301143.
 44. J. Carr, “Neutrino telescopes,” *Nucl. Phys. Proc. Suppl.* **113**, 26 (2002).
 45. A. Letessier-Selvon, “Auger: A large air shower array and neutrino telescope,” astro-ph/0208526.
 46. J. Ahrens *et al.* (AMANDA Collaboration), “Search for supernova neutrino bursts with the AMANDA detector,” *Astropart. Phys.* **16**, 345 (2002) [astro-ph/0105460].
 47. T. Totani, K. Sato, H. E. Dalhed and J. R. Wilson, “Future detection of supernova neutrino burst and explosion mechanism,” *Astrophys. J.* **496**, 216 (1998) [astro-ph/9710203].
 48. The Supernova Early Warning System (SNEWS) <http://hep.bu.edu/~snnet/>
 49. G. G. Raffelt, “Particle physics from stars,” *Ann. Rev. Nucl. Part. Sci.* **49**, 163 (1999) [hep-ph/9903472].
 50. J. F. Beacom, R. N. Boyd and A. Mezzacappa, “Black hole formation in core collapse supernovae and time-of-flight measurements of the neutrino masses,” *Phys. Rev. D* **63**, 073011 (2001) [astro-ph/0010398].

51. A. S. Dighe and A. Y. Smirnov, "Identifying the neutrino mass spectrum from the neutrino burst from a supernova," *Phys. Rev. D* **62**, 033007 (2000) [hep-ph/9907423].
52. C. Lunardini and A. Y. Smirnov, "Probing the neutrino mass hierarchy and the 13-mixing with supernovae," hep-ph/0302033.
53. K. Takahashi and K. Sato, "Effects of neutrino oscillation on supernova neutrino: Inverted mass hierarchy," hep-ph/0205070.
54. K. Takahashi and K. Sato, "Earth effects on supernova neutrinos and their implications for neutrino parameters," *Phys. Rev. D* **66**, 033006 (2002) [hep-ph/0110105].
55. G. G. Raffelt, "Mu- and tau-neutrino spectra formation in supernovae," *Astrophys. J.* **561**, 890 (2001).
56. R. Buras, H. T. Janka, M. T. Keil, G. G. Raffelt and M. Rampp, "Electron-neutrino pair annihilation: A new source for muon and tau neutrinos in supernovae," *Astrophys. J.* in press (2003) [astro-ph/0205006].
57. M. T. Keil, G. G. Raffelt and H. T. Janka, "Monte Carlo study of supernova neutrino spectra formation," astro-ph/0208035.
58. M. Malek *et al.* (Super-Kamiokande Collaboration), "Search for supernova relic neutrinos at Super-Kamiokande," hep-ex/0209028.
59. M. Vagins, "Future detection of SN neutrinos," talk at NOON03, Kanazawa, Japan, 10–14 February 2003.